

Name:
Registration number:
Sex:
ID band:
Date of Birth:
Consultant:

Date:

Instructions and advice following an injection to the eye

Procedure: Intravitreal injection of: to the eye.

- The white of the injected eye may be slightly red and uncomfortable for a few days, especially around the injection site. This is normal and you may take your 'painkillers' if you have any discomfort
- You can take the clear plastic shield off your eye when you get home
- Your vision in that eye may be blurred for several days and you may see 'floaters' in your vision. This is normal and should resolve after a few days
- Please do not get your injected eye wet for 24 hours
- Please do not carry out strenuous physical exercise for 24 hours
- If you have been prescribed post-injection drops, use them as directed

If you experience **any** of the following:

- Increasing redness, swelling or pain in the injected eye
- Loss of vision
- If you are concerned or have any questions

Please contact the Ophthalmology Department:

Area 1, Level 0, Queen Elizabeth Hospital Birmingham,
Mindelsohn Way, Edgbaston, Birmingham B15 2GW.
Telephone: 0121 371 6476 Monday-Friday 9.00am-4.30pm.

Birmingham and Midland Eye Centre: (Out-of-hours)

Sandwell and West Birmingham Hospitals NHS Trust,
Dudley Road, Birmingham B18 7QH
Telephone: 0121 554 3801 ask for Birmingham and Midland Eye Centre