

Home oxygen therapy removal

Following an assessment of your breathing and oxygen levels, it has been decided that you no longer require oxygen therapy at home.

Why don't I need it anymore?

Your recent assessment has shown that oxygen therapy is not beneficial for you at this time. We can tell this from the results of your blood test, called an ear lobe capillary sample, which tell us directly how much oxygen is in your blood. Your breathlessness may be down to your heart or lung condition, not low oxygen levels.

Using oxygen therapy when it is not required can be harmful to your body. In some people it can cause an increase in carbon dioxide levels, which can make you quite unwell, as carbon dioxide is the waste gas that is breathed out from the body. These levels are also measured during your oxygen assessment blood test (as mentioned above). Oxygen is therefore only prescribed to those people who have low oxygen levels and meet oxygen criteria. There may be more effective ways to manage your symptoms.

If I don't need it anymore, why does my breathing feel better when I use oxygen?

You can become psychologically dependent on oxygen but when taught other breathless management techniques (listed below), your confidence can soon return when breathing without it.

Another reason breathing can feel better on oxygen is because you can feel the pressure or flow of oxygen into the nose or mouth. This can make the brain believe you can breathe more easily. Again, this is a psychological, not a physical need for oxygen, as shown by blood tests.

How else can I manage my breathlessness?

We can refer you to the community respiratory therapy service who will give you advice and strategies to help you manage breathlessness, including:

- Breathing techniques
- Inhaler technique
- Body positioning
- Anxiety management
- Using a hand held fan on your face (this can give the same relief as oxygen)
- Pulmonary rehabilitation programme

How will my oxygen therapy be stopped?

You will be weaned off oxygen over a six week period with support if required. The equipment will remain in place over this period and not be removed suddenly. Advice and reassurance will be provided, particularly in managing breathlessness as part of your individual oxygen removal plan. A blood test will be repeated to confirm oxygen therapy is not required.

If you require this information in another format, such as a different language, large print, braille or audio version please ask a member of staff or email patientexperience@uhb.nhs.uk.

Lung Function and Sleep

Good Hope Hospital
Rectory Road, Sutton Coldfield
Birmingham, B75 7RR
Telephone: 0121 371 3870